

490 Hargrave Street Winnipeg, Manitoba R3A OX7 CANADA

www wrha mb ca

490, rue Hargrave Winnipeg (Manitoba) R3A 0X7 CANADA

www wrha mh ca

November 24, 2020

John Taylor Collegiate 470 Hamilton Ave. Winnipeg, MB R2Y 0H4

Dear Parent/Guardian of Students at John Taylor Collegiate,

Manitoba public health officials advised John Taylor Collegiate today of a confirmed case of COVID-19 in the school on November 17-19 when the individual may have been infectious.

John Taylor Collegiate is working closely with public health officials and following their recommendations. Based on the public health investigation, this exposure was assessed to be low risk overall for the school community. Classroom contacts are being advised to monitor for symptoms and seek testing if symptoms develop, effective today. Close contacts of the case have also been identified and advised to self-isolate (quarantine).

Please note: the school is not permitted to release the name or any identifying information about the confirmed case of COVID-19 to any member of the school, community or media.

If you have not been contacted by a letter from public health officials, your child(ren) have not been identified as being exposed to COVID-19 from this case. If your child develops a new onset or worsening of any one symptom listed in Column A or any two or more symptoms listed in Column B (see table below), even if they are mild, immediately isolate them from others. It is recommended your child get tested for COVID-19 as soon as symptoms appear. You may go directly to the nearest COVID-19 testing location during their open hours. For the testing site nearest you, please visit https://www.gov.mb.ca/covid19/updates/testing.html, or call Health Links—Info Santé at 204-788-8200 or toll-free at 1-888-315-9257.

A	В
Fever/Chills	Runny nose
Cough	Muscle aches
Sore throat/hoarse voice	Fatigue
Shortness of breath/difficulty breathing	Pink eye (conjunctivitis)
Loss of taste or smell	Headache
Vomiting or diarrhea for more than 24 hours	Skin rash of unknown cause
	Nausea or loss of appetite
	Poor feeding (if an infant)

Please note: individuals with symptoms outlined in columns A and B, who choose not to be tested, must isolate for a minimum of 10 days following the start of symptoms. Isolation can be stopped at day 10 if the individual has been symptom free for 24 hours.

For COVID-19, public health information about please visit more https://manitoba.ca/covid19/updates/resources.html.

Sincerely,

Dr. Heejune Chang Medical Officer of Health Winnipeg Regional Health Authority