
Open File OF2008-11

CAMIRO Project 97E-02, Thompson Nickel Belt: final report March 2002, revised and updated 2003

by O.M. Burnham, N. Halden, D. Layton-Matthews, C.M. Leshner, J. Liwanag, L. Heaman, L. Hulbert, N. Machado, D. Michalak, M. Pacey, D.C. Peck, A. Potrel, P. Theyer, K. Toope and H. Zwanzig

Winnipeg, 2009

Published by the Manitoba Geological Survey on behalf of the Canadian Mining Industry Research Organization (CAMIRO)

Recommended reference:

Burnham, O.M., Halden, N., Layton-Matthews, D., Leshner, C.M., Liwanag, J., Heaman, L., Hulbert, L., Machado, N., Michalak, D., Pacey, M., Peck, D.C., Potrel, A., Theyer, P., Toohe, K. and Zwanzig, H. 2009: CAMIRO Project 97E-02, Thompson Nickel Belt: final report March 2002, revised and updated 2003; Manitoba Science, Technology, Energy and Mines, Manitoba Geological Survey, Open File OF2008-11, 434 p. + appendices + GIS shape files for use with ArcInfo[®].

Published by:

Manitoba Science, Technology, Energy and Mines
Manitoba Geological Survey
360-1395 Ellice Avenue
Winnipeg, Manitoba
R3G 3P2 Canada

Telephone: (800) 223-5215 (General Enquiry)
(204) 945-4154 (Publication Sales)

Fax: (204) 945-8427

E-mail: minesinfo@gov.mb.ca

Website: manitoba.ca/minerals

Cover photo: Thompson Mine, Thompson, Manitoba

FOREWORD

This report represents a final summary report for CAMIRO Project 97E-02. It includes most of the information in previous reports and summarizes all of the interpretations for the project. It was written by Marcus Burnham, Norm Halden, Larry Heaman, Daniel Layton-Matthews, Michael Leshner, Janice Liwanag, Nuno Machado, Dianne Michalak, David Peck, Alain Potrel, Kim Toope, and Herman Zwanzig. It was compiled by Marcus Burnham, formatted with assistance from Kimberly Nelson and Jo Duke, and edited by Michael Leshner. Kevin Ansdell, Chris Böhm, Carl Chandler, Eric Ducharme, Chris Freund, Denis Gapais, Larry Hulbert, Reid Keays, Jürgen Kraus, Josef Macek, Peter Theyer, and Curtis Wegleitner assisted with field work or contributed samples, data, maps, and/or ideas at various stages of the project. Mark Pacey, Paul Lenton, and John Broome were responsible for compiling the GIS products.

ACKNOWLEDGEMENTS

We are very grateful to the sponsors of this project, Billiton Ltd., Cominco Ltd., Falconbridge Ltd., Hudson Bay Exploration and Development Ltd., Inco Ltd., and WMC Ltd., for their strong and enthusiastic scientific, logistical, and financial support of the project and for giving us this opportunity to work on such interesting and challenging problems. We are especially grateful to Larry Larson, John Lutz, Haddie Mahoney, Scott Mooney, Dave Owens, John Pearson, Jamie Robertson, Rick Somerville, Gary Sorensen, Rob Stewart, Patti Tirschmann, and Kevin Wells for logistical assistance, geological insights, and sampling advice. We benefited greatly from discussions with Wouter Bleeker and Paul Golightly. We are especially grateful to Josef Macek for his long-term contributions to the geology and stratigraphy of the Thompson Nickel Belt and for providing a critical geological and stratigraphic framework for this project. Dick Alcock and Roger Wallis helped us view things from an industry perspective. We are most grateful to David Peck for his skilful coordination of the project during the critical first three years, and to Dick Alcock and Tom Lane for their unflagging support and encouragement through all stages of the project.

TABLE OF CONTENTS

Foreword	iii
Acknowledgements	iii
Table of Contents	iv
List of Figures	xvi
List of Tables	xxvii
Chapter 1 Executive Summary	1.1
1.1 Objectives	1.1
1.2 Results	1.2
1.3 Exploration Guides in TNB	1.4
1.4 Work Not Completed and Follow-Up Work	1.5
Chapter 2 Introduction	2.1
2.1 Preamble	2.1
2.2 Objectives	2.1
2.3 Research Team	2.2
2.4 Sponsors	2.3
2.5 Methodology	2.3
2.5.1 Geology Subproject	2.4
2.5.2 Geochemistry Subproject	2.4
Chapter 3 Regional Geology	3.1
Chapter 4 New Mapping	4.1
4.1 Compilation Maps: Preliminary Release	4.1
4.2 Setting and Mystery Lakes	4.1
4.3 Thompson South Pit	4.2
4.4 Future Mapping	4.2

Chapter 5	Stratigraphy	5.1
5.1	Introduction	5.1
5.2	Stratigraphic Descriptions	5.6
5.2.1	Basement Gneiss (Unit Ag)	5.6
5.2.2	Ospwagan Group (Unit O)	5.6
5.2.2.1	Manasan Formation (Unit Om)	5.8
5.2.2.2	Thompson Formation (Unit Ot)	5.8
5.2.2.3	Pipe Formation (Unit Op)	5.9
5.2.2.4	Setting Formation (Unit Os)	5.10
5.2.2.5	Bah Lake Assemblage (Unit Ob)	5.11
	<i>Setting Lake – Bah Lake Area</i>	5.13
	<i>Upper Ospwagan Lake</i>	5.14
	<i>Mystery Lake</i>	5.15
	<i>Synthesis</i>	5.15
5.2.3	Grass River Group (Unit G)	5.16
5.2.4	Burntwood Group (Unit B)	5.19
5.2.5	Winnipegosis Komatiite Belt	5.19
5.3	Regional Lithofacies and Tectonostratigraphic Interpretation	5.20
5.4	Ultramafic Bodies	5.23
5.4.1	Introduction	5.23
5.4.2	Contact Relationships	5.23
5.4.3	Internal Stratigraphy	5.24
5.4.4	Stratigraphic Location	5.25
5.4.5	Interpretation	5.28
5.4.5.1	Sills vs. Flows	5.28
5.4.5.2	Level of Sill Emplacement	5.28
5.5	Mafic Sills	5.30
5.5.1	Grass River Intrusion	5.31
5.5.2	Soab Intrusion	5.32
5.5.3	Taylor River Intrusion	5.32
5.5.4	Bah Lake Intrusion	5.32
5.5.5	Thompson Mine (South Pit) Intrusion	5.32
5.5.6	Other Intrusions	5.33
5.6	Stratigraphic Interpretations	5.33
5.6.1	Basin Evolution	5.33
5.6.2	Stratigraphic Relationships Between Mafic and Ultramafic Rocks in the TNB	5.38

5.6.3	Molson Dyke Event	5.39
Chapter 6	Structure	6.1
6.0	Preamble	6.1
Part I. Modified Nappe Model		6.1
6.1	Methods and Conventions	6.1
6.2	Structural History	6.2
6.2.1	Previous and Present Work	6.2
6.2.2	Basement Structures (Archean)	6.2
6.2.3	Pre-Grass River Group Proterozoic Structures (D ₁)	6.4
6.2.4	Early Synmetamorphic Structures (D ₂)	6.5
6.2.5	Late Peak-Metamorphic Structures (D ₃)	6.7
6.2.6	Post–Peak Metamorphic Structures (D ₄)	6.8
6.3	Structural geometry and strain patterns	6.9
6.4	Folds	6.10
6.4.1	Archean Basement Structures	6.10
6.4.2	Basement Domes and Basins	6.10
6.4.3	Thompson Nappe	6.12
6.4.4	Soab Creek Structure	6.13
6.4.5	Setting Antiform	6.15
6.4.6	Bah and Fish Lakes Anticlines	6.15
6.4.7	Five Mile Synform	6.16
6.4.8	Kiski Creek Fold	6.16
6.5	Faults and shear zones	6.16
6.5.1	Grass River Lineament	6.17
6.5.2	Burntwood Mylonite Zone	6.18
6.5.3	Setting Lake Fault	6.18
6.5.4	Superior Boundary Fault Zone	6.20
6.5.5	Fish Creek and Pakwa Lake Shear Zones	6.22
6.6	Structural synopsis: discussion and future work	6.22
6.6.1	Revised Tectonic Model and Future Work	6.24
Part II. Transpressional Model		6.26
6.7	Introduction	6.26
6.8	Methodology and Conventions	6.26
6.8.1	Orientation of the Principal Strain Axes	6.26

6.8.2	Type of Finite Strain Ellipsoid	6.27
6.8.3	Strain Intensity	6.27
6.8.4	Shear Criteria	6.27
6.8.5	Strain History	6.28
6.8.6	P-T Conditions	6.29
6.8.7	Geochronology	6.29
6.9	TNB structural data	6.29
6.9.1	Principal Strain Axes and Foliation Trajectories	6.29
6.9.1.1	Foliations	6.29
6.9.1.2	Lineations	6.31
6.9.2	Strain Type and Strain Intensity	6.31
6.9.3	Shear Zones	6.31
6.9.3.1	Geometry of Arrays of Shear Zones	6.31
6.9.3.2	Syn-kinematic Thermal Conditions	6.33
6.9.4	Relationships Between Deformation and Granite Intrusions	6.33
6.9.5	Earlier Deformations	6.34
6.10	Discussion and Interpretation	6.35
Chapter 7	Metamorphism	7.1
7.1	Introduction	7.1
7.2	Ultramafic Rocks	7.1
7.2.1	Introduction	7.1
7.2.2	Alteration	7.2
7.2.3	Petrography	7.3
7.2.3.1	Regional Alteration	7.4
	<i>Serpentinization</i>	7.4
	<i>Alkali-Carbonate Metasomatism</i>	7.5
7.2.3.2	Localized Metasomatism	7.6
7.2.4	Alteration Geochemistry	7.8
7.2.4.1	Regional Alteration	7.8
	<i>General Trends</i>	7.8
	<i>Major Elements</i>	7.8
	<i>Trace Elements</i>	7.11
7.2.4.2	Localized Metasomatism	7.14
7.2.5	Timing of Alteration	7.15
	<i>Serpentinization</i>	7.15
	<i>Pegmatite-Related Metasomatism</i>	7.16

	<i>Alkali and Carbonate Metasomatism</i>	7.16
7.2.6	Conclusions	7.17
7.3	P-T conditions	7.18
7.3.1	Metasediments	7.18
7.3.1.1	Geothermometry	7.19
7.3.1.2	Geobarometry	7.20
7.3.1.3	Summary of P-T Estimates from Metasedimentary Rocks	7.20
7.3.2	Mafic and Ultramafic Rocks	7.21
7.4	Future work	7.22
Chapter 8	Lithogeochemistry and Geochemistry	8.1
8.1	Introduction	8.1
8.2	Aims	8.1
8.3	Sampling	8.1
8.4	Ultramafic Rocks	8.3
8.4.1	Mineralization Status	8.3
8.4.2	Petrography	8.3
8.4.2.1	Metadunites	8.4
8.4.2.2	Metaperidotites	8.5
8.4.2.3	Metapyroxenites	8.6
8.4.3	Olivine Compositions	8.6
8.4.3.1	Textural Characteristics	8.6
8.4.3.2	Oxide Inclusions	8.7
8.4.3.3	Major and Minor Elements	8.9
8.4.3.4	Sulfide Saturation History of Northern TNB Ultramafic Bodies	8.10
	<i>Fractional Crystallization</i>	8.11
	<i>Batch Equilibration</i>	8.11
	<i>Metamorphic Reequilibration</i>	8.12
8.4.3.5	William Lake Olivines	8.14
	<i>Ni Content of the Parental Magma</i>	8.14
	<i>Equilibrium vs. Fractional Crystallization</i>	8.14
	<i>Assimilation-Fractional Crystallization processes</i>	8.15
	<i>Assimilation of Magmatic Sulfides</i>	8.15
	<i>Increasing Oxygen Fugacity</i>	8.15
8.4.3.6	Interim Summary	8.16
8.4.4	Whole-Rock Compositions	8.17
8.4.4.1	Major and Minor Elements	8.17

	<i>Fractional Crystallization-Assimilation models</i>	8.18
	<i>MELTS Models</i>	8.19
	<i>Major Element Compositions of TNB Mafic and Ultramafic Rocks</i>	8.21
	<i>Pearce Element Ratio Diagrams</i>	8.22
	<i>Minor Element Compositions</i>	8.23
	<i>Interim Summary</i>	8.28
8.4.4.2	Lithophile Trace Elements	8.28
	<i>Incompatible Lithophile Elements</i>	8.28
	<i>Magma Source Enrichment vs. Contamination</i>	8.31
	<i>Interim Summary</i>	8.33
8.4.4.3	Highly Chalcophile Trace Elements (PGE and Au)	8.34
8.4.4.4	Nd Isotope Compositions of Mafic and Ultramafic Rocks	8.35
	<i>Amphibolite-Facies Mafic and Ultramafic Rocks in Central TNB Basalts East of William Lake (Winnipegosis Belt)</i>	8.35
	<i>Further Analyses</i>	8.36
8.4.4.5	Summary	8.37
8.4.5	Discriminant Analysis	8.38
8.4.5.1	Introduction	8.38
8.4.5.2	Application to Ultramafic Bodies of the TNB	8.38
	<i>Choice of Elements</i>	8.39
	<i>Training and Test Data Sets</i>	8.39
8.4.5.3	Results	8.40
	<i>Moak Lake (Mineralized)</i>	8.41
	<i>Mystery Lake (Strongly- to Non-mineralized)</i>	8.42
	<i>Thompson (Mineralized)</i>	8.42
	<i>Mid Lake (Weakly Mineralized)</i>	8.42
	<i>Pipe 1 and 2 (Mineralized)</i>	8.42
	<i>Spur South (Non-Mineralized)</i>	8.42
	<i>Northern Winnipegosis Belt (Non-Mineralized)</i>	8.43
	<i>William Lake (Weakly- to Non-Mineralized)</i>	8.43
	<i>Winnipegosis Belt (Non-Mineralized)</i>	8.43
8.4.5.4	Conclusions	8.44
8.5	Mafic Rocks	8.44
8.5.1	Introduction	8.44
8.5.2	Whole rock Compositions	8.44

8.5.2.1	Major and Minor Elements	8.44
	<i>Chromium</i>	8.45
	<i>Nickel</i>	8.45
8.5.2.2	Trace Elements	8.46
	<i>Group 1</i>	8.46
	<i>Group 2</i>	8.46
	<i>Group 3</i>	8.46
	<i>Group 4</i>	8.46
8.5.3	Interim Summary	8.47
8.6	Metasedimentary Rocks	8.47
8.6.1	Introduction	8.47
8.6.2	Petrography	8.48
8.6.3	Whole Rock Compositions	8.48
8.6.3.1	Major and Minor Elements	8.48
	<i>Northern TNB</i>	8.48
	<i>Central TNB</i>	8.51
8.6.3.2	Trace Elements	8.52
	<i>Semi-pelites and Pelites</i>	8.52
	<i>Iron-formations</i>	8.53
	<i>Grass River Group Sedimentary and Tuffaceous Rocks</i>	8.53
8.6.3.3	Nd Isotope Compositions	8.54
8.6.4	Summary	8.54
8.7	Granitoids and Gneisses	8.56
8.8	Interpretations	8.57
8.8.1	Melting Processes and Sources of Mafic and Ultramafic Magmas	8.57
8.8.1.1	Depth and Degree of Partial Melting	8.57
	<i>Depth of Melting</i>	8.58
	<i>Degree and Temperature of Melting</i>	8.58
	<i>Mantle Source Region</i>	8.59
	<i>Metallogenic Implications</i>	8.59
8.8.2	Volcanic Architecture of Mafic and Ultramafic Rocks	8.60
8.8.3	Assimilation Processes During Emplacement	8.62
8.8.4	Environment of Deposition of Ospwagan Group Metasedimentary Rocks	8.64
8.8.5	Exploration Implications	8.64
8.8.5.1	Olivine Compositions	8.64
8.8.5.2	Major and Minor Element Compositions of Ultramafic Bodies	8.65
8.8.5.3	Trace Element Compositions of Ultramafic Bodies	8.67

8.8.5.4	Highly Chalcophile Element Compositions of Ultramafic Bodies	8.69
8.8.5.5	Discriminant Analyses	8.72
8.8.5.6	Sedimentary Rock Compositions	8.72
8.8.5.7	Volcanic Architecture	8.73
8.8.5.8	Melting Processes	8.73
8.8.5.9	Assimilation Processes	8.74
8.9	Future Work	8.75
Chapter 9	Sulfides	9.1
9.1	Introduction	9.1
9.2	Mineralization Status	9.1
9.3	Ore Types	9.2
9.4	Sulfide Textures	9.3
9.4.1	Textural Classification	9.3
9.4.2	Textural Descriptions	9.3
9.4.3	Textural Variations Between Ore Deposits	9.5
9.4.4	Textural Variations Across Strike in a Deformed Ore Zone	9.6
9.4.5	Textural Variations Along Strike in a Deformed Ore Zone	9.6
9.5	Mineral Chemistry	9.7
9.5.1	Pyrrhotite	9.7
9.5.2	Pentlandite	9.8
9.5.3	Trace Elements	9.9
9.5.4	Variably Deformed Sulfide Minerals	9.10
9.5.5	Metal Diffusion	9.10
9.6	Whole Rock Chemistry	9.12
9.6.1	Normalization to 100% Sulfides	9.12
9.6.2	Ni/Co Ratios	9.13
9.6.3	$\delta^{34}\text{S}$ and S/Se Ratios	9.13
9.6.4	Platinum Group Elements (PGEs)	9.14
9.6.5	Trace Elements	9.14
9.6.6	Comparison with Kambalda and Langmuir	9.16
9.7	Interpretations	9.17
9.7.1	Metamorphic Textures	9.17
9.7.2	Magmatic Processes	9.18
9.7.2.1	Chalcophile Element Composition of Parental Magma	9.18

9.7.2.2	Sulfide-silicate ratio (R-factor) and metal contribution from the country rocks	9.20
	<i>Background</i>	9.20
	<i>Metal mixing models</i>	9.21
9.7.2.3	Fractionation of the sulphide liquid	9.25
9.7.3	Platinum Anomaly in Massive Sulfides	9.26
9.7.3.1	Magmatic controls on the behaviour of PT	9.26
9.7.3.2	Hydrothermal controls on the behaviour of PT	9.26
9.7.3.3	Mineralogical controls on the behaviour of PT	9.27
9.7.3.4	Modification of massive ores	9.28
9.7.3.5	Ni enrichment in Thompson ores and sediments	9.29
9.7.4	Role of Deformation in Metal Transport	9.29
9.7.5	Sulfidation	9.30
9.7.6	Compositional and Mineralogical Haloes	9.31
9.8	Conclusions	9.31
9.9	Further Work	9.32
Chapter 10	U-Pb Geochronology	10.1
10.1	Introduction	10.1
10.2	U-Pb Geochronology of Mafic Magmatism	10.2
10.2.1	Introduction	10.2
10.2.2	Sample selection	10.2
10.2.3	U-Pb Results	10.3
10.2.4	Discussion	10.4
10.3	Geochronology of Detrital Rocks	10.5
10.3.1	Introduction	10.5
10.3.2	Provenance of Ospwagan and Grass River Groups Sediments	10.5
10.3.2.1	Ospwagan Group	10.5
10.3.2.2	Grass River Group	10.6
10.3.3	Defining the Western Boundary of the Southern TNB	10.6
10.4	Timing of Tectonic Events	10.7
10.4.1	Introduction	10.7
10.4.2	Eastern Boundary of the TNB	10.7
	<i>Sample TB 20-52a</i>	10.7
10.4.3	Timing of West-Vergent Tectonics	10.9
	<i>Sample TB 20-51</i>	10.9
	<i>Samples TB 99-42 and TB 20-08</i>	10.10

<i>Sample TB 20-11</i>	10.11
<i>Sample TB 20-87a</i>	10.12
<i>Sample TB 20-69</i>	10.12
10.4.4 Youngest Tectonic Events	10.13
<i>Sample 3000/351 AZ1</i>	10.13
<i>Sample TNB 99-21A</i>	10.14
<i>Sample TNB 99-13d</i>	10.15
10.4.5 Summary	10.16
10.4.6 Conclusions	10.16
10.5 Boundaries of Archean Domains in the TNB	10.17
<i>Sample TB 99-27a</i>	10.17
<i>Sample TNB 99-14</i>	10.17
10.5.1 Summary	10.18
10.6 U-Pb geochronology - Summary and Conclusions	10.18
 Chapter 11 Summary and Discussion	 11.1
11.1 Summary of principle results	11.1
11.2 Geological interpretations	11.3
11.2.1 Environment of depositions of the Ospwagan Group	11.3
11.2.2 Source of Ospwagan Group Sediments	11.3
11.2.3 Timing of emplacement of ultramafic sills into the Ospwagan Group	11.3
11.3 Petrogenetic Interpretations	11.4
11.4 Tectonic Interpretations	11.5
11.4.1 Fold-nappe model	11.5
11.4.2 Transpression model	11.6
11.5 Ore genetic interpretations	11.7
11.5.1.1 Intrusion of ultramafic sills	11.8
11.5.1.2 Folding and stretching during high-grade metamorphism	11.9
11.5.1.3 Late stage ductile-brittle deformation	11.9
11.5.1.4 Final Cooling	11.9
11.6 Applications for exploration in the TNB	11.10
11.7 Applications for exploration in other areas	11.11
11.8 Follow-up work	11.12
 Appendix 1 Analytical Techniques	 A1.1
A1.1 Introduction	A1.1

A1.2 Analytical Methods Used During CAMIRO Project 97E-02	A1.1
A1.2.1 Whole-Rock Analyses	A1.1
A1.2.2 Mineral Analyses	A1.27
A1.2.3 Nd-Sm Isotope Analyses	A1.29
A1.3 Filters for Pre-existing Data	A1.29
 Appendix 2 Sample Details in Geochemical Database	 A2.1
A2.1 Sample Numbers	A2.1
A2.2 Sample Locations	A2.1
A2.3 Mineralization Status	A2.4
A2.4 Lithology	A2.9
A2.5 Lithological Units	A2.15
A2.6 Depth in Hole	A2.15
A2.7 Source of Sample	A2.15
A2.8 Summary of Geochemical Data Available	A2.16
A2.9 Methods of Analysis	A2.16
A2.10 Alternate Methods of Analysis	A2.17
A2.11 Raw vs. Volatile-Free Data	A2.17
 Appendix 3 Electron Microprobe Data Used for P-T Estimates	 A3.1
 Appendix 4 Electron Microprobe Data - Olivines and Chromites	 A4.1
 Appendix 5 Electron Microprobe Data - Sulfides	 A5.1
 Appendix 6 PIXE Data - Sulfides	 A6.1
 Appendix 7 Factor Analysis	 A7.1
 Appendix 8 Detrital Zircon $^{207}\text{Pb}/^{206}\text{Pb}$ Ages	 A8.1
 Appendix 9 Felsic Rock Geochemistry	 A9.1
 Appendix 10 U-Pb Ages	 A10.1
 Appendix 11 References	 A11.1

LIST OF FIGURES

Chapter 1 Executive Summary

No Figures

Chapter 2 Introduction

No Figures

Chapter 3 Regional Geology

Figure 3.1	Geological setting of the Thompson Ni Belt.	3.6
Figure 3.2	Bouguer gravity anomaly map of central Manitoba.	3.7

Chapter 4 New Mapping

Figure 4.1	Simplified geology of the southern half of the exposed Thompson Ni Belt.	4.3
Figure 4.2	Simplified geology of the northern half of the exposed Thompson Ni Belt.	4.4

Chapter 5 Stratigraphy

Figure 5.1	Reconstructed Ospwagan Group lithostratigraphy at Thompson pit and Pipe II pit.	5.40
Figure 5.2	Simplified stratigraphic sections of the Ospwagan Group in the central TNB.	5.41
Figure 5.3	Tectonostratigraphic sections of the Setting-August lakes area.	5.42
Figure 5.4	Schematic interpretation of tectonostratigraphic and intrusive relationships in the TNB and adjacent part of the Kisseynew Domain.	5.43
Figure 5.5	Geological interpretation for the Winnipegosis Komatiite Belt.	5.44
Figure 5.6	Graphical logs summarizing the geology of typical ultramafic bodies from the northern and central TNB.	5.45
Figure 5.7	Graphical logs summarizing the geology of ultramafic bodies containing amphibolite layers from the central TNB.	5.46
Figure 5.8	Variation in petrology of ultramafic and/or amphibolite sills as a function of differentiation and crystal settling.	5.47
Figure 5.9	Areas used to group the ultramafic bodies of the Northern TNB during spatial analysis.	5.48
Figure 5.10	Pie charts of the distribution of host lithologies of the TNB ultramafic bodies.	5.49
Figure 5.11	Stratigraphic index for ultramafic bodies of the TNB as a function of location.	5.50
Figure 5.12	Models for the control of sediment density on the depth of sill emplacement.	5.51

Chapter 6 Structure

Figure 6.1	Schematic longitudinal sections through proposed fold-nappes.	6.37
Figure 6.2	Schematic cross-sections through the regional structures.	6.38
Figure 6.3	Simplified geological map, structural interpretation, and equal-area stereograms of southern Setting Lake.	6.39

Figure 6.4	Simplified map, structural interpretation, and stereograms of northern Setting Lake.	6.40
Figure 6.5	Equal-area stereograms of the fabric elements in the northern Setting Lake area.	6.41
Figure 6.6	Equal area stereograms of the fabric elements in the southern Setting and Pakwa Lake areas.	6.42
Figure 6.7	Equal-area stereograms of the fabric elements west of the Setting Lake area.	6.43
Figure 6.8	Equal-area stereograms of the fabric elements north of the Setting Lake area	6.44
Figure 6.9	Equal-area stereograms of fabric elements and structures related to mylonite, shear zones, and faults in the Setting Lake and surrounding areas.	6.45
Figure 6.10	Cartoons illustrating the interpreted structural evolution of the Oswagan, Grass River, and Burntwood Groups in the northern Setting Lake area.	6.46
Figure 6.11	Interpretation of structural relationships at the south end of Setting Lake.	6.47
Figure 6.12	Geophysical and geological interpretations of the seismic reflection profile crossing the Superior Boundary near Ponton.	6.48
Figure 6.13	Geophysical and geological interpretations of the seismic reflection profile crossing the Superior Boundary near Easterville.	6.49
Figure 6.14	Transformation of a sphere during pure and simple shear.	6.50
Figure 6.15	Orientation of structures relative to principal strain axes during flattening strain.	6.50
Figure 6.16	Different types of finite strain ellipsoids.	6.51
Figure 6.17	Experimentally produced faults nets for coaxial and non-coaxial deformation.	6.51
Figure 6.18a	Flinn diagram showing expected geometry of shear zone arrays.	6.52
Figure 6.18b	Theoretical effect of bulk shearing deformation on the distribution of shear directions and shear senses on individual shear zones.	6.52
Figure 6.19	Example of progressive deformation and the development of C/S/C' structures.	6.53
Figure 6.20	General foliation map of the TNB and adjacent eastern domain.	6.54
Figure 6.21a	Oswagan Lake foliation trajectories and lineations map.	6.55
Figure 6.21b	Setting Lake foliation trajectories map.	6.56
Figure 6.22	Stereograms showing attitudes of regional stretching lineation and foliation.	6.57
Figure 6.23	Photographs of field locations showing:	
	a) Top to the west thrust in amphibolite facies conditions, South Jonas Road.	6.58
	b) Reverse top to the W fault, Thompson T3 Pit.	6.58
	c) Dextral mineralized shear zone under greenschist facies conditions, Thompson T3 Pit.	6.58
	d) Melt-bearing dextral shear bands in the Archean gneisses, Thompson T3 Pit.	6.59
	e) Folded boudins in a pegmatitic vein intrusive into Oswagan Group sediments, Thompson T3 Pit.	6.59
	f) Vertical stretching on mylonitic schistosity.	6.59
	g) Post-folial pegmatite associated with top to the W motion intrusive in the Archean gneisses.	6.59
Figure 6.24	Stereograms and frequency histograms illustrating the distribution of shear zones and shear bands in the exposed TNB.	6.60

Figure 6.25	Frequency histograms showing distribution of measured shear zones according to their strikes and kinematics.	6.61
Figure 6.26	Cartoon illustrating the structural pattern observed in the TNB.	6.62

Chapter 7 Metamorphism

Figure 7.1	Metamorphic map of the Thompson – Pipe area of the TNB.	7.24
Figure 7.2	Photomicrographs of pseudomorph and transitional styles of serpentinization in the TNB ultramafic bodies.	7.25
Figure 7.3	Photomicrographs of non-pseudomorph and vein-style serpentinization in the TNB ultramafic bodies.	7.26
Figure 7.4	Photomicrographs of alkali and carbonate alteration of TNB ultramafic bodies.	7.27
Figure 7.5	Schematic diagrams showing the mobility of elements during the production of alteration zones in ultramafic rocks adjacent to granitic pegmatites.	7.28
Figure 7.6	Photomicrographs of alteration zones adjacent to granitic pegmatites.	7.29
Figure 7.7	Variation in LOI as a function of MgO and CO ₂ content for mafic and ultramafic rocks from the TNB.	7.31
Figure 7.8	Mantle-normalized multi-element plot showing elemental enrichment accompanying K metasomatism.	7.32
Figure 7.9	Plot of [La/Sm] _n vs. total LOI for serpentinized TNB ultramafic bodies.	7.32
Figure 7.10	Primitive mantle normalized REE plot, showing the high La/Ce ratios of some samples from the TNB.	7.33
Figure 7.11	Mantle-normalized La/Ce ratios as a function of LOI, K/K*, CO ₂ , Eu/Eu* and H ₂ O for ultramafic rocks from the central TNB.	7.34
Figure 7.12	Plot of [La/Yb] _{mn} vs. [Nb/Yb] _{mn} for TNB ultramafic rocks.	7.35
Figure 7.13	Trace element enrichment as a function of Yb and MgO content in the Mystery Lake ultramafic body (DDH 89225, 89234 and 89236).	7.36
Figure 7.14	Enrichment-depletion diagrams showing compositional changes between alteration zones adjacent to granitic pegmatites.	7.37
Figure 7.15	Photomicrographs of metapelites and metabasalts used in P-T estimates for the central TNB.	7.38
Figure 7.16	Location of the drill holes and samples used in geothermometry and geobarometry.	7.39
Figure 7.17	Summary of geothermometric and geobarometric data obtained for metasedimentary rocks during this study.	7.40

Chapter 8 Silicate Rock Mineralogy and Geochemistry

Figure 8.1	Photomicrographs of textures and mineralogy of olivine adcumulate rocks from the TNB.	8.76
Figure 8.2	Photomicrographs of textures and mineralogy of olivine orthocumulate and pyroxenitic rocks from the TNB.	8.79
Figure 8.3	Compositions of chromites and opaque inclusions in olivines from the cores of ultramafic bodies from the TNB.	8.80
Figure 8.4	Histogram of relict magmatic olivine compositions from TNB ultramafic bodies.	8.81

Figure 8.5	Equilibrium and average olivine compositional variations with fractional crystallization of a komatiitic magma.	8.81
Figure 8.6	MnO contents of TNB olivines plotted as a function of forsterite content.	8.82
Figure 8.7	Ni contents of TNB olivines plotted as a function of forsterite content.	8.83
Figure 8.8	Comparison of measured olivine compositions with those expected for fractional crystallization of olivine and sulfide from a sulfide-saturated magma.	8.84
Figure 8.9	Comparison of measured olivine compositions with those expected for batch equilibration of a evolving magma with an externally derived sulfide liquid.	8.85
Figure 8.10	A model for the development of high Ni contents in the olivines of the W-56 ultramafic body.	8.86
Figure 8.11	Major element compositions of liquids and cumulate assemblages produced during pure fractional crystallization of an ultramafic magma.	8.87
Figure 8.12	Major, compatible minor, and incompatible, immobile trace element contents of mafic and ultramafic rocks from the northern TNB plotted as a function of MgO.	8.88
Figure 8.13	Major, compatible minor, and incompatible, immobile trace element contents of mafic and ultramafic rocks from the central TNB plotted as a function of MgO.	8.93
Figure 8.14	Major, compatible minor, and incompatible, immobile trace element contents of mafic and ultramafic rocks from the southern TNB plotted as a function of MgO.	8.96
Figure 8.15	Pearce element ratio plots of molar (Mg + Fe + Mn)/Ti vs. Si/Ti for S-poor samples from TNB ultramafic bodies.	8.99
Figure 8.16	Pearce element ratio plots of molar Mg/Ti vs. (Fe+Mn)/Ti for S-poor mafic and ultramafic rocks from the central TNB.	8.100
Figure 8.17	Comparison of typical trace element patterns for a komatiitic magma and upper crustal sediments.	8.101
Figure 8.18	[La/Sm] _{mn} vs. [Th/Nb] _{mn} for ultramafic bodies of the TNB.	8.102
Figure 8.19	[Th/Nb] _{mn} vs. MgO for mafic and ultramafic rocks from the TNB.	8.103
Figure 8.20	[Th/Nb] _{mn} vs. MgO for mafic and ultramafic rocks from the TNB.	8.104
Figure 8.21	Condensed mantle normalized multi-element plot illustrating the expected trace element variations expected for variable degrees of melt-depletion, source enrichment and contamination during the production and emplacement of ultramafic magmas.	8.105
Figure 8.22	Mantle-normalized Nb/Ti ratios as a function of Th/Ti for different mixing and/or contamination processes.	8.106
Figure 8.23	Mantle-normalized Nb/Ti vs. Th/Ti for ultramafic rocks from the TNB.	8.107
Figure 8.24	Variation in highly chalcophile element abundances as a function of MgO content for mafic and ultramafic rocks from all regions of the TNB.	8.109
Figure 8.25	ϵ_{Nd} vs. time for mafic and ultramafic rocks from the Pipe 1 ultramafic body and the central TNB.	8.110
Figure 8.26	$^{143}Nd/^{144}Nd$ plotted as a function of incompatible element enrichment.	8.111
Figure 8.27	Discriminant function plot showing data for all TNB ultramafic bodies.	8.112
Figure 8.28	Discriminant function plots for individual TNB ultramafic bodies.	8.113
Figure 8.29	Komatiite-tholeiite discriminant plot for mafic and ultramafic volcanic rocks and associated sills and dykes in the TNB and WKB.	8.115

Figure 8.30	Jensen cation plot for mafic and ultramafic volcanic rocks and associated sills and dykes in the TNB and WKB.	8.116
Figure 8.31	Whole-rock Cr and MgO contents for mafic and ultramafic volcanic rocks and associated dykes and sills in the TNB and WKB.	8.117
Figure 8.32	Whole-rock MgO and Ni contents for mafic and ultramafic volcanic rocks and associated sills and dykes in the TNB and WKB.	8.118
Figure 8.33	Mantle normalized Ce/Sm and Nb/Ti ratios for mafic and ultramafic rocks in the TNB and WKB.	8.119
Figure 8.34	Mantle normalized Ce/Sm and Th/Nb ratios for mafic and ultramafic rocks in the TNB and WKB.	8.119
Figure 8.35	Mantle-normalized trace element abundances for selected, representative samples of mafic and ultramafic volcanic rocks and associated dykes and sills in the TNB and WKB.	8.120
Figure 8.36	Chemostratigraphy of Oswagan Group metasedimentary rocks from the type section at Pipe Pit.	8.121
Figure 8.37	Chemical classification of TNB metasedimentary rocks and basement gneisses from the type sections at Pipe and Thompson Pits using $\log(\text{Fe}_2\text{O}_3/\text{K}_2\text{O})$ vs. $\log(\text{SiO}_2/\text{Al}_2\text{O}_3)$.	8.122
Figure 8.38	Discriminant function diagram for the sandstone-mudstone suites from basement paragneisses and metasedimentary rocks at Pipe and Thompson Pits.	8.122
Figure 8.39	Cr vs. Zr in metasedimentary rocks and basement gneisses from the type sections at Pipe and Thompson Pits.	8.123
Figure 8.40	Chemical classification of metasedimentary rocks from the central TNB using $\log(\text{Fe}_2\text{O}_3/\text{K}_2\text{O})$ vs. $\log(\text{SiO}_2/\text{Al}_2\text{O}_3)$.	8.123
Figure 8.41	Discriminant function diagram for sandstone-mudstone suites from the central TNB, based on major element compositions.	8.124
Figure 8.42	Cr vs. Zr in metasedimentary rocks from the central TNB.	8.124
Figure 8.43	Multi-elemental plots for Oswagan Group metasedimentary rocks from the type sections at the Pipe and Thompson Pits.	8.125
Figure 8.44	Mantle-normalized HFSE compositions of metapelites, quartzites, iron formations and basement rocks of the northern and central TNB.	8.126
Figure 8.45	Multi-elemental plots for Oswagan Group iron formations from the type sections at the Pipe and Thompson Pits.	8.126
Figure 8.46	Co vs. P_2O_5 plot for the Pipe Formation iron formations relative to the average compositions of other iron formations within and around the Superior Province.	8.127
Figure 8.47	Mantle-normalized plots of metasediments and tuffs of the Grass River Group.	8.128
Figure 8.48	ϵ_{Nd} vs. time for sedimentary rocks from the TNB.	8.129
Figure 8.49	Classification of the felsic rocks of the TNB according to their molecular normative An-Ab-Or compositions.	8.129
Figure 8.50	A/NK versus A/CNK diagram for the felsic magmatism of the TNB.	8.130
Figure 8.51	Plot of A/CNK versus SiO_2 illustrating the calc-alkaline affinity of the felsic magmatism of the TNB.	8.130
Figure 8.52	Comparison of SiO_2 contents in experimental melts obtained at a range of pressures with the inferred compositions of the parental magmas to the Bah Lake volcanic rocks and TNB ultramafic bodies.	8.131

Figure 8.53	MgO contents of experimental melts as functions of the degree of melting and temperature.	8.132
Figure 8.54	Three possible models for the timing of emplacement of the ultramafic sills in the Ospwagan Group of the Thompson Ni Belt.	8.133
Figure 8.55	Schematic, interpretive model showing inferred petrogenetic relationships.	8.134
Figure 8.56a	Schematic, interpretive model showing decoupling of lithophile and chalcophile component in and the thermal aureole around <i>mineralized undifferentiated cumulate feeder sills</i> in the Ospwagan Group of the Thompson Nickel Belt.	8.135
Figure 8.56b	Schematic, interpretive model showing decoupling of lithophile and chalcophile component in and the thermal aureole around <i>non-mineralized undifferentiated cumulate feeder sills</i> in the Ospwagan Group of the Thompson Nickel Belt.	8.136
Figure 8.56c	Schematic, interpretive model showing decoupling of lithophile and chalcophile component in and the thermal aureole around <i>non-mineralized differentiated non-cumulate sills</i> in the Ospwagan Group of the Thompson Nickel Belt.	8.137

Chapter 9 Sulfide Mineralogy and Geochemistry

Figure 9.1	Textures representing disseminated sulfides in metasedimentary rocks from the TNB.	9.33
Figure 9.2	Textures representing disseminated sulfides in ultramafic rocks and interstitial sulfides in ultramafic rocks from the TNB.	9.34
Figure 9.3	Textures representing barren sulfides in metasedimentary rocks from the TNB.	9.35
Figure 9.4	Textures representing massive sulfides in ultramafic breccias from the TNB.	9.36
Figure 9.5	Textures representing massive and semi-massive sulfides in metasedimentary rocks from the TNB.	9.37
Figure 9.6	BSE image showing semi-massive sulfides hosted in biotite gneiss from the 1D ore body, Thompson mine.	9.38
Figure 9.7	Brecciated silicate fragments along contact between garnet biotite schist and massive sulfides.	9.39
Figure 9.8	Stringer sulfides in garnet biotite gneiss from the Thompson T1 Mine.	9.39
Figure 9.9	Example of chalcopyrite occurring adjacent to porphyroblastic silicates in massive sulfides.	9.40
Figure 9.10	Pyrite-chalcopyrite symplectites.	9.41
Figure 9.11	Massive sulfides from the Thompson T1 Mine.	9.42
Figure 9.12	Massive sulfides from the Thompson 1C ore body.	9.43
Figure 9.13	Semi-massive sulfides from the Thompson 1D ore body.	9.44
Figure 9.14	Schematic diagram of an ore zone in the 1C ore body, Thompson Mine.	9.45

Figure 9.15	Reflected light photomicrographs of typical sulfide-bearing samples from a Ni ore zone in the 1C ore body, Thompson Mine.	9.46
Figure 9.16	Relative concentrations of Fe, Ni, and Co in pentlandite from a range of sulfide types from the Thompson 1C ore body.	9.47
Figure 9.17	Relative concentrations of Fe, Ni, and S in pentlandite from a range of sulfide types from the Thompson 1C ore body.	9.48
Figure 9.18	Relative concentrations of Fe, Ni, and S in pyrrhotite from a range of sulfide types from the Thompson 1C ore body.	9.49
Figure 9.19	Electron microprobe point analyses across an equant porphyroblastic pentlandite grain and adjacent pyrrhotite in two dimensions.	9.50
Figure 9.20	Concentration of Ni in co-existing sulfide minerals from samples along section I-I' in Figure 9.14, Thompson 1C ore body.	9.51
Figure 9.21	Concentration of Ni in pyrrhotite from sulfide-bearing samples along section J-J' in Figure 9.14, Thompson 1C ore body.	9.52
Figure 9.22	Ni and S concentrations in monoclinic pyrrhotite from the Thompson Mine (T1 mine, 1C ore body, and 1D ore body).	9.53
Figure 9.23	Ni and S concentrations in monoclinic pyrrhotite from massive and semi-massive sulfides in metasedimentary rocks and semi-massive sulfides in Archean Basement (Bucko only).	9.53
Figure 9.24	Ni and S concentrations in monoclinic pyrrhotite from massive sulfides in ultramafic breccias.	9.54
Figure 9.25	Ni and S concentrations in monoclinic and hexagonal pyrrhotite from disseminated sulfides in ultramafic rocks.	9.54
Figure 9.26	Ni, S, and Ni/Co in pentlandite from massive and semi-massive sulfides in metasedimentary rocks and semi-massive sulfides in Archean Basement (Bucko only).	9.55
Figure 9.27	Ni, S, and Ni/Co in pentlandite from massive sulfides in ultramafic breccias.	9.56
Figure 9.28	Ni, S, and Ni/S in pentlandite from disseminated sulfides in ultramafic rocks.	9.57
Figure 9.29	Ni and Se in monoclinic pyrrhotite from massive and semi-massive sulfides in metasedimentary rocks.	9.58
Figure 9.30	Ni and Se in pentlandite from massive and semi-massive sulfides in metasedimentary rocks.	9.58
Figure 9.31	Ni in massive and semi-massive sulfides from the TNB, normalized to 100% sulfides.	9.59
Figure 9.32	Ni/Co ratios of ultramafic bodies from the TNB.	9.59
Figure 9.33	Ni/Co ratios of massive and semi-massive sulfides from the TNB.	9.60

Figure 9.34	Se/S and $\delta^{34}\text{S}$ data for Ni sulfide ore and barren sedimentary sulfides from the TNB.	9.60
Figure 9.35	Se/S ratios of massive and semi-massive sulfide samples from the TNB.	9.61
Figure 9.36	Se/S of ultramafic bodies from the TNB.	9.61
Figure 9.37	Chondrite-normalized plot of chalcophile element concentrations in massive and semi-massive sulfide samples from the TNB.	9.62
Figure 9.38	Chondrite-normalized plot of chalcophile element concentrations in samples from Birchtree Mine.	9.62
Figure 9.39	Chondrite-normalized plot of chalcophile element concentrations in samples from Thompson Mine.	9.63
Figure 9.40	Chondrite-normalized plot of chalcophile element concentrations in samples from the Pipe deposit.	9.63
Figure 9.41	Pd and Bi concentrations in massive and semi-massive sulfide samples from the TNB.	9.64
Figure 9.42	Bismuthotellurides in massive sulfides from the 1D ore body, Thompson mine.	9.65
Figure 9.43	Ni and As concentrations in massive and semi-massive sulfides from the TNB.	9.66
Figure 9.44	Ni and Pd concentrations in massive and semi-massive sulfides from the TNB.	9.66
Figure 9.45	Ni and Bi concentrations in massive and semi-massive sulfides from the TNB.	9.67
Figure 9.46	Chalcophile element compositions of parental magmas to TNB ultramafic bodies.	9.67
Figure 9.47	Compositions of magmatic sulfides produced for different R factors and metal contents in the external sulfide liquid.	9.68
Figure 9.48	Solubility of Pd and Pt in the hydrous silicate melt <i>versus</i> oxygen fugacity.	9.70
Figure 9.49	Mantle-normalized chalcophile element patterns for ores from the TNB, Kambalda, Langmuir, Donaldson West, Lac des Iles, New Rambler, Rathbun Lake, Kuroko, and Besshi.	9.71
Figure 9.50	Schematic model showing theoretical variations in metal concentration profiles for physical mixing, fluid infiltration, and solid-state diffusion.	9.74
Figure 9.51	Schematic geochemical dispersion model for an ore deposit in the TNB that has been deformed transpressively.	9.75
Figure 9.52	Schematic model showing variations in Au/Cu/Pt dispersion accompanying variations in deformation.	9.75

Chapter 10 U-Pb Geochronology

Figure 10.1	Locations of mafic dykes sampled for U-Pb age dating during CAMIRO Project 97E-02.	10.20
Figure 10.2	Compilation of ages for mafic and ultramafic samples from the TNB.	10.21

Figure 10.3	Histograms of $^{207}\text{Pb}/^{206}\text{Pb}$ ages of detrital zircons from the Oswagan Group.	10.22
Figure 10.4	Comparison of age distributions for the Oswagan Group and the western Superior Province.	10.23
Figure 10.5	Histograms of age distributions for detrital zircons of the Grass River, Burntwood, Missi and Sickle Groups.	10.24
Figure 10.6	$^{207}\text{Pb}/^{206}\text{Pb}$ age distribution of detrital zircons from the Oswagan and Grass River Groups.	10.25
Figure 10.7	$^{207}\text{Pb}/^{206}\text{Pb}$ age distribution of detrital zircons from five drill holes drilled on Falconbridge's William Lake, Bracken Lake, and South Moose Lake properties.	10.26
Figure 10.8	Tectonic model for the TNB proposed by Bleeker (1990b) and modified by White et al. (1999).	10.27
Figure 10.9	Sampling locations for U-Pb study of felsic rocks and pegmatites.	10.28
Figure 10.10	Concordia diagram for zircon and monazite from pegmatite sample TB 20-52a.	10.29
Figure 10.11	Concordia diagram for zircon from granite sample TB 20-51.	10.30
Figure 10.12	Concordia diagram for zircon and monazite from granite samples TB 99-42 and TB 20-08.	10.30
Figure 10.13	Concordia diagram for zircon and monazite from pegmatite samples TB 20-11 and TB 20-87a.	10.31
Figure 10.14	Concordia diagram for zircon from granite sample TB 20-69.	10.32
Figure 10.15	Concordia diagram for zircon and monazite from pegmatite sample 3000/351 AZ1.	10.32
Figure 10.16	Concordia diagram for zircon and monazite from pegmatite sample TB 99-21a.	10.33
Figure 10.17	Concordia diagram for monazite from pegmatite sample TB 99-13d.	10.34
Figure 10.18	Concordia diagram for zircon from Archean gneiss sample TB 99-14.	10.34
Figure 10.19	Plot of TNB U-Pb metamorphic ages as a function of distance from the Kisseynew Domain.	10.35
Figure 10.20	U-Pb metamorphic ages plotted relative to generalized foliation trends for the TNB.	10.36

Chapter 11 Summary and Discussion

No Figures

Appendix 1 Analytical Techniques

Figure A1.1	Precision of major element fusion XRF data determined from analyses of in-house standards and duplicate samples.	A1.3
Figure A1.2	Precision of minor element pressed pellet XRF data determined from analyses of in-house standards and duplicate samples.	A1.6
Figure A1.3	Precision of major and minor element concentrations determined by ICP-AES analyses of in-house standards, duplicate samples and reference materials.	A1.9
Figure A1.4	Accuracy of major and minor element ICP-AES data for sulfide-rich samples determined from analyses of reference materials.	A1.12
Figure A1.5	Comparison of Ni and Cu concentrations determined by pressed-pellet XRF, ICP-AES and flame-AAS for sulfide-rich samples.	A1.14
Figure A1.6	Comparison of Cr concentrations determined by INAA and XRF.	A1.15

Figure A1.7	Precision of ICP-MS analyses of selected elements using replicate analyses of in-house standards (n > 3).	A1.16
Figure A1.8	Precision of total C and S determinations by IR-spectrometry.	A1.19
Figure A1.9	Accuracy of S data by infrared spectroscopy determined from analyses of reference materials.	A1.20
Figure A1.10	95% confidence intervals for Se, Sb and Bi obtained by hydride-generation AAS.	A1.22
Figure A1.11	Comparison of Se data by hydride-generator atomic absorption spectrometry (HG-AAS) and instrumental neutron activation (INAA).	A1.23
Figure A1.12	Precious metal concentrations of CANMET standards WPR-1 and TDB-1 by NiSFA-ICP-MS.	A1.24

Appendix 2 Sample Details in Geochemical Database

Figure A2.1	Locations of sampling areas for geochemical samples in the northern TNB.	A2.2
Figure A2.2	Locations of sampling areas for geochemical samples in central and southern TNB.	A2.3
Figure A2.3	Summary of mineralization status of sampling locations in northern TNB.	A2.7
Figure A2.4	Summary of mineralization status of sampling locations in southern and central TNB.	A2.8

Appendix 3 Electron Microprobe Data Used for P-T Estimates

No Figures

Appendix 4 Electron Microprobe Data - Olivines and Chromites

No Figures

Appendix 5 Electron Microprobe Data - Sulfides

No Figures

Appendix 6 PIXE Data - Sulfides

No Figures

Appendix 7 Factor Analysis

No Figures

Appendix 8 Detrital Zircon – $^{207}\text{Pb}/^{206}\text{Pb}$ Ages

No Figures

Appendix 9 Felsic Rock Geochemistry

No Figures

Appendix 10 U-Pb Ages

No Figures

Appendix 11 References

No Figures

LIST OF TABLES

Chapter 1 Executive Summary

No Tables

Chapter 2 Introduction

No Tables

Chapter 3 Regional Geology

No Tables

Chapter 4 New Mapping

No Tables

Chapter 5 Stratigraphy

Table 5.1	TNB stratigraphic units used in maps and text	5.2
Table 5.2a	TNB stratigraphy (Central and southernmost part of the northern region)	5.4
Table 5.2b	TNB stratigraphy (Central and northern region)	5.5
Table 5.3	Densities used in modelling of ultramafic sill emplacement	5.29

Chapter 6 Structure

Table 6.1	Summary of dominant TNB structure	6.3
-----------	-----------------------------------	-----

Chapter 7 Metamorphism

Table 7.1	Primary and alteration mineral assemblages present in TNB ultramafic bodies	7.3
Table 7.2	Results of principal component analysis of whole rock major, minor, and trace element contents of ultramafic rocks from the TNB	7.10
Table 7.3	Results of principal component analysis of trace element ratios in ultramafic rocks from the TNB	7.13
Table 7.4	Metamorphic mineral assemblage in metapelitic rocks from the central TNB	7.18
Table 7.5	Summary of geobarometric and geothermometric calculations for metasedimentary rocks from the central TNB	7.20
Table 7.6	Summary of metamorphic mineral assemblages from mafic and ultramafic rocks from the central TNB	7.21
Table 7.7	Summary of geothermometric calculations for mafic rocks from the central TNB	7.22

Chapter 8 Silicate Rock Mineralogy and Geochemistry

Table 8.1	Summary of samples in project database	8.2
Table 8.2	Whole-rock compositions of components used in MELTS models	8.20

Table 8.3	Consequences of contamination on the Si and Fe contents, oxidation state, and chromite and sulfide saturation states of komatiitic magmas	8.25
Table 8.4	Nd isotope compositions of mafic and ultramafic samples	8.36
Table 8.5	Mineralisation discriminant function coefficients	8.40
Table 8.6	Classification matrix for TNB ultramafic bodies	8.41
Table 8.7	Descriptions of metasedimentary samples collected from central region of TNB	8.50
Table 8.8	Nd isotope compositions of metasedimentary rocks	8.55
Table 8.9	Nd isotope compositions of Archean gneisses	8.56
Table 8.10	Amount of sulfides produced, ore tenor, ore and magma geochemical characteristics in terms of flow duration and flow rate for a high-Mg komatiite system	8.70

Chapter 9 Sulfide Mineralogy and Geochemistry

Table 9.1	Sulfide ore types, textures, distributions, and host rocks in the TNB	9.3
Table 9.2	Compositions of sulfide minerals in samples from the Thompson 1C ore body	9.11
Table 9.3	Chalcophile element and PGE data for massive and semi-massive sulfides in the TNB	9.15
Table 9.4	Trace element data for massive and semi-massive sulfides from the TNB	9.16
Table 9.5	Composition of magma used in sulfide modelling	9.19
Table 9.6	Mass-conservation equations for two-component silicate-sulfide systems	9.21
Table 9.7	Parameters used in modeling the magmatic sulfide compositions	9.22

Chapter 10 U-Pb Geochronology

Table 10.1	Summary of Mafic Samples Collected from TNB	10.2
------------	---	------

Chapter 11 Interpretations and Models

No Tables

Appendix 1 Analytical Techniques

Table A1.1	Major element analyses of in-house reference materials by fusion XRF	A1.2
Table A1.2	Summary of analytical reproducibility of major element analyses by fusion XRF	A1.3
Table A1.3	Major element compositions of reference materials determined by fusion XRF	A1.4
Table A1.4	Minor element analyses of in-house reference materials by powder pellet XRF	A1.5
Table A1.5	Summary of the reproducibility of minor elements by pressed pellet XRF	A1.6
Table A1.6	Minor element analyses of reference materials determined by pressed pellet XRF	A1.7
Table A1.7	Major and minor element compositions obtained on sulfide-poor samples by ICP-AES	A1.10
Table A1.8	Major and minor element compositions obtained on sulfide-rich samples by ICP-AES	A1.11
Table A1.9	Calculated corrections for ICP-AES analyses	A1.13
Table A1.10	Comparison of Cu and Ni data obtained by ICP-AES, PP-XRF, and flame-AAS	A1.13
Table A1.11	Cr data for SRM analysed by INAA	A1.14

Table A1.12	Trace element data for in-house reference materials analysed by ICP-MS	A1.16
Table A1.13	Regression of trace element data for reference materials analysed by ICP-MS	A1.17
Table A1.14	Total C and S concentrations determined for reference materials by IR-spectrometry	A1.20
Table A1.15	As and Se data for reference materials analysed by INAA	A1.22
Table A1.16	Precious metal concentrations determined for CANMET reference materials WPR-1 and TDB-1 by Ni-FA ICP-MS	A1.25
Table A1.17	Priorities given to sources of data for the compilation of the geochemical database	A1.26
Table A1.18	EMPA conditions and data quality during the analysis of silicate minerals	A1.27
Table A1.19	Data for in-house silicate and oxide mineral standards analysed by electron microprobe	A1.28

Appendix 2 Sample Details in Geochemical Database

Table A2.1	Mineralization state of ultramafic bodies in the northern TNB	A2.4
Table A2.2	Mineralization state of ultramafic bodies in the central and southern TNB	A2.6
Table A2.3	Lithological classification scheme used in geochemical database	A2.9
Table A2.4	Mapping between the lithological codes used in the contributed databases and codes used within the project geochemical database	A2.10
Table A2.5	Codes used to indicate the data available for each sample	A2.16
Table A2.6	Method codes used in geochemical database	A2.16

Appendix 3 Electron Microprobe Data Used for P-T Estimates

Table A3.1	Electron microprobe data used in the calculation of pressures and temperatures from metasedimentary samples	A3.2
Table A3.2	Electron microprobe data used in the calculation of temperatures from mafic samples	A3.4

Appendix 4 Electron Microprobe Data - Olivines and Chromites

Table A4.1	Compositions of olivines from ultramafic bodies of the TNB	A4.2
Table A4.2	Compositions of cumulus chromite and oxide inclusions from ultramafic bodies of the TNB	A4.7

Appendix 5 Electron Microprobe Data - Sulfides

Table A5.1	Electron Microprobe Data – Sulfides	A5.2
------------	-------------------------------------	------

Appendix 6 PIXE Data - Sulfides

Table A6.1	Major and trace element data obtained by PIXE for TNB sulfides	A6.2
------------	--	------

Appendix 7 Factor Analysis

No Tables

Appendix 8 Detrital Zircon $^{207}\text{Pb}/^{206}\text{Pb}$ Ages

Table A8.1	$^{207}\text{Pb}/^{206}\text{Pb}$ ages of detrital zircons from the Ospwagan Group	A8.1
------------	--	------

Table A8.2	$^{207}\text{Pb}/^{206}\text{Pb}$ ages for detrital zircons from the Grass River Group	A8.4
Table A8.3	$^{207}\text{Pb}/^{206}\text{Pb}$ ages for detrital zircons from drill holes on Falconbridge's William Lake, Bracken Lake, and South Moose Lake permits	A8.5

Appendix 9 Felsic Rock Geochemistry

Table A9.1	Major and trace element data for granitoid and gneiss samples	A9.1
Table A9.2	Major element data for granitoid and gneiss samples collected in 2000	A9.2

Appendix 10 U-Pb Ages

Table A10.1	U-Pb data for timing of tectonic events	A10.1
-------------	---	-------

Appendix 11 References

No Tables